

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

NOGALLTM

ACTIVE CONSTITUENT: 1,000 million Agrobacterium radiobacter var. K1026/g peat

A bacterial culture for the control of crown gall disease on stone fruit trees, nut trees and roses.

Net Contents: 100g, 250g

BASF Australia Ltd ABN 62 008 437 867
Level 12, 28 Freshwater Place Southbank VICTORIA 3006, AUSTRALIA
Website: www.agro.basf.com.au

Customer Service Hotline: 1800 558 399

APVMA Approval No: 39304/100g/0905, 39304/250g/0905



DIRECTIONS FOR USE

CROP	DISEASE	RATE	CRITICAL COMMENTS
Stone Fruit Almonds Walnuts Pecans Roses	Crown Gall	250g per 12L water or 100g per 5L water	Seeds: Dip and thoroughly wet seeds before planting. Seedlings: Dip and thoroughly wet roots within 2 hours of lifting. Trim damaged roots before dipping. Cuttings: Dip before applying rooting compounds. However, when using alcohol based solutions, dip cuttings in accordance with directions, allow to dry for 2 hours then treat with NOGALL™

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS:

NOGALL™ is a bacterial culture, contained in a peat medium, which controls crown gall disease (*Agrobacterium tumefaciens*) by the inoculation of seeds, seedlings and cuttings. It is a living culture which must be protected from extremes of temperature and direct sunlight. Plant material should be dipped at each transfer stage, if possible.

Mixing

Place one packet of NOGALL™ into a clean container, add the required quantity of cool, clean water and stir vigorously. Keep thoroughly stirred during use. Use the solution within 24 hours.

Compatibility

DO NOT use with any other pesticides, fertilizers or chemicals which may harm the bacteria.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the bacteria culture or used container.

STORAGE AND DISPOSAL

Store in the closed original container in a dry, well-ventilated area as cool as possible, out of direct sunlight. Dispose of empty packaging by wrapping in plastic bag and placing in garbage or local authority landfill. If any dip solution is excess, add approximately 50ml household bleach per 1L, stir thoroughly and allow to react for approximately 4 hours prior to disposal.

SAFETY DIRECTIONS

Wash hands after use. Avoid breathing dust.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 1126).

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the manufacturer.

WARRANTY

The product must be used strictly in accordance with the label directions and instructions. BASF accepts no liability or responsibility for any consequential loss or otherwise arising from failure to follow these directions and instructions. The liability of BASF is limited to that imposed by statutes and, in any case, the liability is strictly limited to replacement of the product or (at the option of BASF) the refund of the price paid.

APVMA Approval No: 39304/100g/0905, 39304/250g/0905

Batch No:

Date of Manufacture:

NOGALL

LABEL



BASF Australia Ltd
ABN 62 008 437 867
Level 12, 28 Freshwater Place
Southbank VICTORIA 3006
AUSTRALIA

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE – ALL HOURS – AUSTRALIA WIDE