

Inoculant Group Chart

Nodulaid[®] Inoculant

Noculator Inoculant Granules

**Please refer to Row Spacing table on back for amount required

С	All Subterranean, Crimson, Rose, Cupped & Helmet Clovers Arrowleaf, Balansa, Purple and Persian Clover		√
	Pea, Field Pea, Faba Bean and Vetch Seed	500 kg	
E/F	Lentil	250 kg	✓
\mathbf{c}	Lupin Seed and Serradella Pods	500 kg	/
G	Yellow, Pink, Hybrid & Slender Serradella Seeds	250 kg	
Ν	Chickpea	500 kg	√
		Maximum weight of seeds treated (kg) per 1.25 kg pack	

*Nodulaid Inoculants E, F, G & N contain Celstik



Inoculant row spacing

Use the appropriate application rate based on this table:

Row spacing (inches)	Row spacing (cm)	Application rate (kg/ha)
7	17.8	6
8	20.3	5.3
9	23	4.6
10	25.4	4.2
11	27.9	3.8
12	30.5	3.5
13	33	3.2
14	36.6	3
15	38.1	2.8
16	40.6	2.6

Apply granular inoculant directly to the furrow. The product must not be applied at a depth that is less than the planting depth of the seed.

- Do not mix Nodulator with granular herbicides and fertilisers during planting.
- Environmental conditions may affect Nodulator's flowability. Check the metering system regularly to ensure proper flow.
- Nodulator and Nodulaid contain living organisms. Always use each pack before its expiry date.



For more information on BASF inoculants, visit **crop-solutions.basf.com.au** or contact your local BASF representative on **1800 558 399**

