Voraxor Herbicide

THE SIGN OF NEW CONFIDENCE SIMPLER EARLY-SEASON BROADLEAF

WEED CONTROL WITH ARG ASSIST

Growers can now achieve simpler, more comprehensive pre-sowing weed control. Voraxor[®] provides both knockdown and long-lasting pre-emergent residual control of broadleaf weeds, which will help delay the need for post-emergent sprays. Voraxor also has additive impact on annual ryegrass that can strengthen management of cereal cropping's most damaging weed.



What will Voraxor add to my weed control program?

Very versatile early-season weed control, including up to 12 weeks' residual control of key broadleaf weeds – filling a big gap in previous pre-emergent weed control programs.



What's the easiest and most cost-effective way to use Voraxor?

At 200 mL/ha in a tank-mix with a pre-emergent grass weed specialist for one-pass residual control with enhanced activity on ARG. For knockdown control, add glyphosate or paraguat and 1% Hasten to the mix.



How is Voraxor applied and incorporated?

In the same way as the other modern pre-emergent cereal herbicides: applied by boom spray and incorporated by sowing ensuring good physical separation between the seed and treated soil.



Is Voraxor a knockdown or pre-emergent herbicide?

Voraxor is both. As a pre-emergent herbicide, Voraxor will give growers up to 12 weeks' residual control of key BLWs and will also knockdown up to 29 species of emerged broadleaf weeds.

If glyphosate or paraguat is included in the tank-mixture, Voraxor will extend its knockdown spectrum and enhance the burndown control of annual ryegrass - including glyphosateresistant biotypes.





• How does Voraxor perform on the most hard-to-control BLWs?

These percentages are the averaged results 200 mL/ha of Voraxor produced across five years of development trials:

Wild radish **87%** Fleabane **94%** Prickly lettuce **100%** Wireweed **94%** Sowthistle/milk thistle **92%** Capeweed **88%**

Composite data from all development trials in APVMA submission 2013–17. Voraxor applied at 200 mL/ha (0-7 days) and incorporated by sowing (IBS). Control assessed at 56–115 DAT.



A

A

Will Voraxor save me a post-emergent herbicide spray?

It really depends on your paddock history and the resulting weed seed bank. What it should do every year is reduce the pressure to get into the paddock early. You will have more flexibility in the timing of in-crop sprays and more modes of action to choose from.



A

How will Voraxor help me manage annual ryegrass?

Voraxor has proven activity against annual ryegrass, so unless you're already achieving 100% ARG control it should always provide some additional impact. Voraxor is registered to knock down ARG – including glyphosate-resistant seedlings – when it's tank-mixed with either glyphosate or paraquat. If you're using it in paddocks with more than 50% sand in the top 10cm, Voraxor will also suppress ARG for up to 12 weeks and enhance the results achieved by pre-emergent ARG products like Luximax[®] and Sakura^{*}.



Three use patterns

200 mL/ha 0–7 days before IBS

- ✓ Residual control
- Knockdown control (+ 1% Hasten^{*})

240 mL/ha 7–21 days before IBS

- ✓ Residual control
- Knockdown control (+ 1% Hasten^{*})

3 100 mL/ha + 1% Hasten Fallow + pre-plant

✓ Knockdown only

Key attributes

Mode of action group:

PPO Inhibitors (Group 14, formerly Group G)

Crop registrations:

Wheat, barley, durum, oats (grain, hay, forage), triticale

Weed registrations:

Residual control: 16 broadleaf weeds Residual suppression of annual ryegrass in light textured soils

Knockdown control:

29 broadleaf weeds standalone Enhanced control of ARG, including glyphosate-resistant biotypes, in tank-mixes with glyphosate or paraquat.

Find out more about **Voraxor**°**Herbicide** at **crop-solutions.basf.com.au** or call **1800 558 399**

ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

This leaflet is intended as general advice. The information submitted in this publication is based on current BASF knowledge and experience. In view of the many factors that may affect its application, this data does not relieve the user from carrying out their own tests. The data does not imply assurance of certain properties or of suitability for a specific purpose. It is the responsibility of the user to ensure that any proprietary rights and existing laws and legislation are observed. © Copyright BASF 2022 ® Registered trademark of BASF * Registered trademark. 212537 09.2022

