

# Poncho® Plus

Insecticidal Seed Treatment

FACT SHEET

## Poncho® Plus Insecticidal Seed Treatment in pasture and forage brassicas

### Highly productive pest protection for emerging seedlings

Poncho® Plus offers a broad spectrum of pest control in pasture that has been shown to increase the quantity and quality of feed. It combines two powerful active ingredients that complement each other to protect crops and pasture against a range of the most damaging pests for up to four weeks after sowing.



- Excellent control of cutworm in pastures.
- Protection from cockchafer and African black beetle damage for up to four weeks after sowing.
- Allows grass pastures to be cut or grazed 6 weeks after sowing.
- Helps produce higher quantities of fodder with high ME.

#### Application rate

500 mL/100 kg seed

#### WHPs

#### Harvest

Not required when used as directed.

#### Grazing and cutting for feed

Grass pasture: 6 weeks.

Forage brassicas and broadleaf pasture: 8 weeks.



### Pasture and feed crop registrations

- Control
- Suppression

	Forage brassicas	Broadleaf pasture	Grass pasture
Aphids	■ Control		
Cutworms	■ Control	■ Control	■ Control
Wireworms	■ Control		
Blue oat mites	■ Control	■ Control	■ Control
Red-legged earth mites	■ Control	■ Control	■ Control
Yellow-headed cockchafer			■ Control
African black beetles			■ Control
Lucerne fleas	■ Suppression	■ Suppression	■ Suppression


**BASF**  
We create chemistry

# Poncho® Plus

## Insecticidal Seed Treatment

### Cutworm control


Mean mortality in forage crops & pasture



Averaged results from 8 trials: two each in forage brassicas, medica and phalaris; one each in ryegrass and subclover. Trial IDs: WO8-226,W10-353,W10-354.

### Improved productivity

Dry matter production



PGG Wrightson trial at Terang, VIC

### Complementary mite control

Poncho Plus will protect emerging seedlings from mite damage for 3–4 weeks after sowing. Monitoring should start within this period to determine the need for supplementary control measures.

Use Poncho Plus as part of an integrated mite management program which may include:

#### For autumn sowing

After a pasture phase, sow seed treated with Poncho Plus following a well-timed spring insecticide spray (prior to the development of diapause eggs). After a cropping phase, a spring insecticide spray is not usually required unless monitoring in spring finds moderate mite populations.

#### For spring sowing

At the end of a pasture phase, monitor for mite activity and if necessary apply an insecticide spray prior to sowing seed treated with Poncho Plus.

### Resistance management

Both of the active ingredients in Poncho Plus belong to the neocotinoid class of insecticides and Poncho Plus is classified as a **Group 4A** insecticide. When additional control measures are required, avoid repeated use of insecticides with the same mode of action.

### Protection of bees and other insect pollinators

Poncho Plus is highly toxic to bees and should only be used as a seed treatment. DO NOT perform seeding operations under very dry or windy conditions to avoid exposing pollinators to treated seed dust.



For more information on Poncho® Plus, scan this QR code or contact your local BASF representative on **1800 558 399**



#### ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FACT SHEET.

This fact sheet is intended as general advice. Disclaimer: The information submitted in this publication is based on current BASF knowledge and experience. In view of the many factors that may affect its application, this data does not relieve the user from carrying out their own tests. The data does not imply assurance of certain properties or of suitability for a specific purpose. It is the responsibility of the user to ensure that any proprietary rights and existing laws and legislation are observed.