

Voraxor[®]

Herbicide



VICTORY OVER MORE WEEDS

ONE-PASS PREMIUM KNOCKDOWN & PRE-EM BROADLEAF & GRASS WEED CONTROL



NOW REGISTERED
in 8 crops incl. select pulses

Growers can now achieve simpler, more comprehensive pre-sowing weed control. Voraxor[®] provides both knockdown and long-lasting pre-emergent residual control of broadleaf weeds, which will help delay the need for post-emergent sprays. Voraxor also has additive impact on annual ryegrass that can strengthen management of cereal cropping's most damaging weed.

Q What will Voraxor add to my weed control program?

A Very versatile early-season weed control, including up to 12 weeks' residual control of key broadleaf weeds – filling a big gap in previous pre-emergent weed control programs.

Q What's the easiest and most cost-effective way to use Voraxor?

A At 200 mL/ha in a tank-mix with a pre-emergent grass weed specialist for one-pass residual control with enhanced activity on ARG. Application at this timing with 1% high-quality MSO added to the mix also provides fast, reliable knockdown. Trials have shown that Voraxor makes a very effective substitute for paraquat as the follow-up spray in a double knock.

Q How is Voraxor applied and incorporated?

A In the same way as the other modern pre-emergent cereal herbicides: applied by boom spray and incorporated by sowing ensuring good physical separation between the seed and treated soil.

Q Is Voraxor a knockdown or pre-emergent herbicide?

A Voraxor is both. As a pre-emergent herbicide, Voraxor will give growers up to 12 weeks' residual control of key BLWs and will also knockdown up to 29 species of emerged broadleaf weeds.

If glyphosate or paraquat is included in the tank-mixture, Voraxor will extend their knockdown spectrum and enhance the burndown control of annual ryegrass – including glyphosate-resistant biotypes.

BASF
We create chemistry

Q How does Voraxor perform on the most hard-to-control BLWs?

A These percentages are the averaged results 200 mL/ha of Voraxor produced across five years of development trials:

Wild radish 87%	Wireweed 94%
Fleabane 94%	Sowthistle/milk thistle 92%
Prickly lettuce 100%	Capeweed 88%

Composite data from all development trials in APVMA submission 2013–17. Voraxor applied at 200 mL/ha (0–7 days) and incorporated by sowing (IBS). Control assessed at 56–115 DAT.

Q Will Voraxor save me a post-emergent herbicide spray?

A It really depends on your paddock history and the resulting weed seed bank. What it should do every year is reduce the pressure to get into the paddock early. You will have more flexibility in the timing of in-crop sprays and more modes of action to choose from.

Q How will Voraxor help me manage annual ryegrass?

A Voraxor has proven activity against annual ryegrass, so unless you're already achieving 100% ARG control it should always provide some additional impact. Voraxor is registered to knock down ARG – including glyphosate-resistant seedlings – when it's tank-mixed with either glyphosate or paraquat. If you're using it in paddocks with more than 50% sand in the top 10cm, Voraxor will also suppress ARG for up to 12 weeks and enhance the results achieved by pre-emergent ARG products like Luximax® and Sakura*.

Q Can Voraxor be used throughout my crop rotation?

A Not quite. We recommend you don't use the same chemistry in the paddock every year anyway, but Voraxor is now registered in all major cereal crops and chickpeas, faba beans and field peas. That allows you to use different tank-mixes with grass herbicides to ensure you can maintain high-performing weed control through almost every phase.

WEED smart
every weed every seed
every farm every year



Find out more about
Voraxor® Herbicide
at crop-solutions.basf.com.au
or call **1800 558 399**

ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

This leaflet is intended as general advice. The information submitted in this publication is based on current BASF knowledge and experience. In view of the many factors that may affect its application, this data does not relieve the user from carrying out their own tests. The data does not imply assurance of certain properties or of suitability for a specific purpose. It is the responsibility of the user to ensure that any proprietary rights and existing laws and legislation are observed. © Copyright BASF 2024 ® Registered trademark of BASF * Registered trademark. AU55-P00000211 11224

VORAXOR IN A SNAPSHOT

Three use patterns

- 1 200 mL/ha
0–7 days before IBS**
 - ✓ Residual control
 - ✓ Knockdown control
(+ 1% high-quality MSO)
- 2 240 mL/ha
7–21 days before IBS**
 - ✓ Residual control
 - ✓ Knockdown control
(+ 1% high-quality MSO)
- 3 100 mL/ha
+ 1% high-quality MSO
Fallow + pre-plant**
 - ✓ Knockdown only

Key attributes

Mode of action group:

PPO Inhibitors
(Group 14, formerly Group G)

Crop registrations:

Wheat, barley, durum, oats
(grain, hay, forage), triticale,
chickpeas, faba beans, field peas

Weed registrations:

Residual control:

16 broadleaf weeds

Residual suppression of annual ryegrass
in light textured soils

Knockdown control:

29 broadleaf weeds standalone

Enhanced control of ARG, including
glyphosate-resistant biotypes, in
tank-mixes with glyphosate or paraquat.

BASF
We create chemistry