

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Liberty<sup>®</sup> ULTRA

Herbicide

ACTIVE CONSTITUENT: 211 g/L GLUFOSINATE-P-AMMONIUM

GROUP	<b>10</b>	HERBICIDE
-------	-----------	-----------

For non-residual control of certain broadleaf and grass weeds in LibertyLink<sup>®</sup> Canola varieties (with tolerance to Liberty<sup>®</sup> Ultra Herbicide), fallow and optical spot spray technologies (OSST), as specified in the DIRECTIONS FOR USE table.

**IMPORTANT: READ THE SAFETY DIRECTIONS BEFORE USING THIS PRODUCT**

CONTENTS: 1L - 1000L

**BASF Australia Ltd**  
ABN 62 008 437 867  
Level 23, 40 City Road, Southbank VICTORIA 3006  
[crop-solutions.basf.com.au](http://crop-solutions.basf.com.au)

® Registered trademark of BASF

APVMA Approval No.: 94738/143503



## RESTRAINTS

- DO NOT apply to canola varieties other than LibertyLink® Canola varieties with tolerance to Liberty® Ultra Herbicide.
- DO NOT apply by aircraft
- DO NOT apply by mister
- DO NOT apply when rain is expected within 4 hours or final weed control may be reduced
- DO NOT apply if heavy rains or storms are forecast within 3 days.
- DO NOT irrigate to the point of water runoff from the treatment area for at least 3 days after application.
- DO NOT apply after the early bolting stage (BBCH 33) in canola.
- DO NOT apply onto weeds when dew, fog or mist is present.
- DO NOT apply to weeds under stress due to, for example, very dry, very wet, frosty, windblasted insect damaged, nutrient deficient or diseased conditions.

## SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets not smaller than a MEDIUM spray droplet size category
- minimum distances between the application site and downwind sensitive areas (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

### Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones (metres)				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 2000 mL/ha	0.5 m or lower	0	10	0	0	0
	1.0 m or lower	10	40	0	20	0
Up to 1600 mL/ha	0.5 m or lower	0	10	0	0	0
	1.0 m or lower	10	30	0	15	0
Up to 1100 mL/ha	0.5 m or lower	0	0	0	0	0
	1.0 m or lower	5	25	0	10	0
800 mL/ha or lower	0.5 m or lower	0	0	0	0	0
	1.0 m or lower	0	20	0	10	0

DO NOT apply by Optical Spot Spray Technologies (OSST) unless the following requirements are met:  
 -spray droplets not smaller than a MEDIUM spray droplet size category  
 -the release height is not greater than 0.8 metres above the ground

### Buffer zones for Optical Spot Spray Technologies (OSST)

Application rate	Mandatory downwind buffer zones (metres)				
	Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to a maximum of 2000 mL/ha	0	30	0	10	0

### DIRECTIONS FOR USE

TABLE 1 - LIBERTYLINK® CANOLA VARIETIES

CROP	WEED	WEED STAGE	RATE	CRITICAL COMMENTS	
<b>USE ONLY WITH LIBERTYLINK® CANOLA VARIETIES (WITH TOLERANCE TO LIBERTY® ULTRA HERBICIDE ).</b>					
<b>LibertyLink® Canola varieties (with tolerance to Liberty® Ultra Herbicide)</b>  Apply Liberty® Ultra Herbicide to crop from cotyledons fully expanded through to early bolting stage.	Annual ryegrass ( <i>Lolium rigidum</i> )	2 to 4 leaf/start of tillering (Z12 to Z21)	1.1 L/ha followed by 1.1 L/ha 7 - 14 days later <b>OR</b> 1.6 L/ha followed by 1.6 L/ha 7 - 14 days later	Apply Liberty® Ultra Herbicide to crop from cotyledons fully expanded through to early bolting stage.  <b>IT IS ESSENTIAL TO COMPLETE BOTH APPLICATIONS.</b>  The sequential application is designed for a weed control program.	
	Barley grass ( <i>Hordeum leporinum</i> ), Brome grass ( <i>Bromus</i> spp.)		0.8 L/ha followed by 0.8 L/ha 7 - 14 days later	Liberty® Ultra Herbicide does not provide residual weed control. Apply the second Liberty® Ultra Herbicide application 7 to 14 days after the first Liberty® Ultra Herbicide application. A shorter interval between applications will result in better weed control. Complete coverage of weeds is essential for good control. Poor coverage may result in re-growth.	
	Capeweed ( <i>Arctotheca calendula</i> )	2 to 8 leaf			
	Chickpea ( <i>Cicer arietinum</i> ), Corn gromwell ( <i>Lithospermum arvense</i> ), Doublegee/Spiny emex/ Three cornered jack ( <i>Emex australis</i> ), Field pea ( <i>Pisum sativum</i> ), Fumitory ( <i>Fumaria</i> spp.), Lentil ( <i>Lens culinaris</i> ), Loosestrife ( <i>Lythrum hyssopifolia</i> ), Lupin/narrowleaf lupin ( <i>Lupinus angustifolius</i> ), Medic ( <i>Medicago</i> spp.), Prickly lettuce ( <i>Lactuca serriola</i> ), Scotch thistle ( <i>Onopordum acanthium</i> ), Silver grass/squirrel-tailed fescue ( <i>Vulpia myuros</i> )	2 to 4 leaf			<b>Annual ryegrass</b> Use the higher rate of Liberty® Ultra Herbicide where the weed populations are high or growing conditions are less than ideal.  For optimal annual ryegrass control in <b>Liberty Tolerant LibertyLink® Canola varieties</b> ; applications of Liberty® Ultra Herbicide should be used in conjunction with pre- and early post-emergence applications of suitable registered herbicides e.g. pre-emergence applications of metazachlor, propyzamide or trifluralin and post-

<b>LibertyLink® Canola varieties (with tolerance to Liberty® Ultra Herbicide)</b>  Apply Liberty® Ultra Herbicide to crop from cotyledons fully expanded through to early bolting stage.	Crassula/Stone crop ( <i>Crassula</i> spp.)	2 to 8 leaf	0.8 L/ha followed by 0.8 L/ha 7 - 14 days later	emergence applications of clethodim on susceptible annual ryegrass.  <b>Wild radish</b> Do not apply to dense populations of wild radish (>25 weeds/m <sup>2</sup> ).  <b>Seek advice from BASF Australia Ltd before using products other than Liberty® Ultra Herbicide on Dual Tolerant LibertyLink® Canola varieties</b>
	Subterranean clover ( <i>Trifolium subterraneum</i> ), Toad rush ( <i>Juncus bufonius</i> ), Tree hogweed ( <i>Polygonum patulum</i> ), Vetch ( <i>Vicia sativa</i> )	2 to 4 leaf		
	Volunteer barley ( <i>Hordeum vulgare</i> ), Volunteer oats ( <i>Avena sativa</i> ), Volunteer wheat ( <i>Triticum aestivum</i> )	2 leaf to 3 tillers (Z12 to Z23)		
	Wild oats ( <i>Avena</i> spp.)	2 leaf to 2 tillers (Z12 to Z22)		
	Wireweed ( <i>Polygonum aviculare</i> )	2 to 4 leaf		
	Yellow burweed/amsinckia ( <i>Amsinckia intermedia</i> )	2 to 6 leaf		
	<b>Suppression only</b> Wild radish ( <i>Raphanus raphanistrum</i> )	2 to 4 leaf	1.6 L/ha followed by 1.6 L/ha 7 to 14 days later	
<b>LibertyLink® Canola varieties (with tolerance to Liberty Ultra Herbicide)</b> grown for seed production only	Non-tolerant canola plants	2 to 4 leaf  4 to 6 leaf  up until time of bolting	2 L/ha followed by 2 L/ha  2 L/ha, if necessary	Two applications are required to remove the segregating wild type plants. A third application, if necessary, may be applied to the canola up until time of bolting. <b>Growers using Liberty® Ultra Herbicide for canola hybrid seed production do so under contract to BASF Australia Ltd or its agent(s).</b>

**APPLICATION OF LIBERTY® ULTRA HERBICIDE TO CANOLA VARIETIES OTHER THAN LIBERTYLINK® CANOLA VARIETIES (WITH TOLERANCE TO LIBERTY® ULTRA HERBICIDE) WILL RESULT IN SEVERE CROP INJURY OR DEATH OF THE CROP.**

**TABLE 2 - SUMMER FALLOW SITUATIONS**

CROP/ SITUATION	WEEDS	WEED STAGE	RATE	CRITICAL COMMENTS
<p>Prior to starting a fallow, fallow maintenance and prior to planting;</p> <p><b>Cereal grains</b> (including wheat, barley, oats, maize and sorghum)</p> <p><b>Pulses</b> (including chickpeas, faba beans, field peas, lentils, lupins and mungbeans),</p> <p><b>Oilseeds</b> (including canola, cotton, soybeans and sunflowers)</p> <p>Do not sow crops until 1 day or more have elapsed after the final application.</p>	<p><b>Control of:</b> Annual ryegrass (<i>Lolium rigidum</i>), Annual polymeria (<i>Polymeria</i> spp.), Bellvine (<i>Ipomoea plebeia</i>), Bladder ketmia (<i>Hibiscus trionum</i>), Caltrop (<i>Tribulus terrestris</i>), Dwarf amaranth (<i>Amaranthus macrocarpus</i>), Field bindweed (European bindweed) (<i>Convol vulus arvensis</i>), Flax-leaf fleabane (<i>Erigeron bonariensis</i>), Paddy melon (<i>Cucumis myriocarpus</i>), Peach vine (<i>Ipomoea lonchop hylla</i>), Red pigweed (<i>Portulaca</i> spp.), Rhyncho (<i>Rhynchosia</i> spp.), Sesbania pea (<i>Sesbania</i> spp.), Sowthistle (Milk thistle) (<i>Sonchus oleraceus</i>), Volunteer cotton (<i>Gossypium hirsutum</i>) (other than glufosinate tolerant cotton), Yellow vine (<i>Tribulus micrococcus</i>)</p> <p><b>Suppression of:</b> Chinese lantern (Wild gooseberry) (<i>Physalis minima</i>), Noogoora burr complex (<i>Xanthium</i> spp.)</p>	<p>Broadleaf weeds: 2 to 6 leaf</p> <p>Grass weeds: pre-tillering</p>	<p>2 L/ ha in a minimum of 100 L water</p>	<p>Apply to actively growing weeds. Good coverage is essential. Refer 'Application' section for details.</p> <p><b>DO NOT</b> apply more than three applications per season.</p> <p>Liberty® Ultra Herbicide will have an effect on weeds that are larger than the recommended leaf stage, but speed of activity and level of control may be reduced.</p> <p><b>CLIMATIC CONDITIONS</b> <b>Best results are achieved when Liberty® Ultra Herbicide is applied under warm humid conditions</b> (temperatures below 33 °C with a relative humidity above 50 %). Under any other conditions efficacy and speed of action may be reduced.</p>

**TABLE 3 - OPTICAL SPOT SPRAY TECHNOLOGIES (OSST)**

Note: Calibrate the sprayer to spray the equivalent of 100 L/ha. DO NOT apply greater than 37 L of spray mixture per hectare through OSST equipment (2 L of product per hectare)

CROP/SITUATION	WEEDS	WEED STAGE	RATE	CRITICAL COMMENTS
<p>Prior to starting a fallow, fallow maintenance and prior to planting;</p> <p><b>Cereal grains</b> (including wheat, barley, oats, maize and sorghum),</p> <p><b>Pulses</b> (including chickpeas, faba beans, field peas, lentils, lupins and mung beans),</p> <p><b>Oilseeds</b> (including canola, cotton, soybeans and sunflowers)</p> <p>Do not sow crops until 1 day or more have elapsed after the final application.</p>	<p>Awnless barnyard grass (<i>Echinochloa colona</i>) Feathertop Rhodes grass (<i>Chloris virgata</i>) Milkthistle/Sowthistle (<i>Sonchus oleraceus</i>) Pigweed (<i>Portulaca oleracea</i>) Stinkgrass (<i>Eragrostis cilianensis</i>) Windmill grass (<i>Chloris truncata</i>)</p>	Budding/seed head emergence – active flowering	4 to 5.4 L/100 L	<p>Apply to actively growing weeds. Good coverage is essential. Refer to Application section for details. Where rate range occurs use higher rate for actively flowering plants</p> <p><b>DO NOT</b> apply more than three applications per season.</p> <p><b>Liberty Ultra</b> will affect weeds that are larger than the recommended leaf stage, but speed of activity and level of control may be reduced.</p> <p><b>CLIMATIC CONDITIONS</b> Best results are achieved when <b>Liberty Ultra</b> is applied under warm humid conditions (temperatures below 33°C with a relative humidity above 50%). Under any other conditions efficacy and speed of action may be reduced.</p>
	<p>Dwarf amaranth (<i>Amaranthus macrocarpus</i>) Bladder ketmia (<i>Hibiscus trionum</i>) Flaxleaf fleabane (<i>Erigeron bonariensis</i>) Liverseed grass (<i>Urochloa panicoides</i>)</p>	Up to active flowering	5.4 L/100 L	
Weeds listed in Table 2	<p>Broadleaf weeds: 2 to 6 leaf</p> <p>Grass weeds: pre-tillering</p>			

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS**

**HARVEST**

NOT REQUIRED WHEN USED AS DIRECTED.

**GRAZING**

**CANOLA:** DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.

**CANOLA GROWN FOR SEED PRODUCTION:** DO NOT GRAZE OR CUT FOR STOCK FEED.

**SUMMER FALLOW:** (including optical spot spray technology): DO NOT GRAZE OR CUT FOR STOCK FOOD A CROP SOWN FOLLOWING A FALLOW SPRAY FOR 6 WEEKS AFTER SOWING.

**All other situations:** DO NOT GRAZE OR CUT TREATED AREAS FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION

## EXPORT OF TREATED PRODUCE

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Liberty® Ultra Herbicide. If you are growing produce for export, please check with BASF Australia Ltd for the latest information on MRLs and import tolerances BEFORE using Liberty® Ultra Herbicide.

## GENERAL INSTRUCTIONS

Liberty® Ultra Herbicide is a non-volatile herbicide with selective activity against many annual and perennial broadleaf weeds and grasses in LibertyLink® Canola varieties that are specially developed to be tolerant to Liberty® Ultra Herbicide. Ensure spray tank is thoroughly decontaminated prior to use of Liberty® Ultra Herbicide to ensure all chemical residues are removed. Apply Liberty Ultra® Herbicide to the LibertyLink® canola from cotyledon to early bolting stage. Liberty® Ultra Herbicide is absorbed by plant foliage and green stems. It is inactive in soil and does not provide residual weed control. Liberty Ultra® Herbicide is not significantly translocated as an active herbicide throughout the plant, and therefore will only kill that part of a green plant that is contacted by spray. The shorter interval between applications will give better weed control. Best results are achieved when applications are made under good growing conditions to young, actively growing weeds. Liberty® Ultra Herbicide will have an effect on weeds that are larger than the recommended leaf stage, but speed of activity and control may be reduced. Application to weeds under stress (e.g. due to continuous severe frosts, dry or waterlogged conditions, windblasted, insect damaged, nutrient deficient or diseased conditions) should be avoided.

## Climatic Conditions

Best results are achieved when Liberty® Ultra Herbicide is applied under mild humid conditions. Poor results may occur under hot, dry conditions. Under cool (below 10°C), dry and low relative humidity conditions speed of action and control may be reduced.

## Symptoms

Visible symptoms of weed control appear in 3 to 7 days, but complete desiccation may take 20 to 30 days under cool conditions. Minor transient crop effects may be visible after application of Liberty® Ultra Herbicide to LibertyLink® Canola varieties (with tolerance to Liberty® Ultra Herbicide).

## Mixing Instructions

Liberty® Ultra Herbicide mixes easily with water. Clean water should always be used for mixing with Liberty® Ultra Herbicide. Ensure that the spray tank is free of any residues of previous spray materials. Maintain agitation whilst filling the tank and throughout the spraying operation.

Liberty® Ultra Herbicide is a soluble concentrate (SL) formulation. When using in a tank mix with other products the following mix order should be observed;

1. Partly fill the spray tank with water, as per label directions, to at least 70% full. **Commence agitation** and maintain agitation during the mixing and spraying process.
2. Add any **Water Conditioners**.
3. Add any **Wettable Powders (WP)** or **Dry Flowable (DF)** products
4. Add any **Water Dispersible Granules (WDG)** products.
5. **Add any Flowable Liquids (FL) or Suspension Concentrates (SC).**
6. Add any **Emulsifiable Concentrates (EC).**
7. Add any **Aqueous Concentrate (AC) or Soluble (Liquid) Concentrates (SL)** such as Liberty® Ultra Herbicide
8. Add **Adjuvant** if required.
9. Fill the remainder of the spray tank with water.

Use the spray mix within 24 hours of preparation (unless otherwise stated on label).

### Use of Surfactant/Wetting agent

The addition of a surfactant is not required with Liberty® Ultra Herbicide.

### Application Equipment

Ground Boom Sprayers Only

Aim to apply a thorough and even coverage of spray to the entire target plant. Dense stands of weeds should be thoroughly wetted with spray. Incomplete coverage due to e.g. stubble retention, crop plants or weed leaf to leaf overlap, may result in poor control. Equipment should be such that adequate coverage, penetration and volume of spray liquid can be achieved.

Liberty® Ultra Herbicide should be applied at the rate shown in the Directions for Use table in sufficient water to give thorough coverage of weeds. It is recommended that 80 to 100 L water/ha is applied with spray droplets of a MEDIUM droplet size category. In some situations (e.g. high weed density) increased water volumes are likely to give higher levels of weed control.

### Sprayer cleanup

Clean all equipment after use by thoroughly flushing with water.

### Compatibility

#### Canola

Liberty® Ultra Herbicide may be mixed with Decis®\* Options, dimethoate, Lontrel Advanced®\*, Nufarm Crucial®\* Select®\*, Select®\* Xtra, Targa®\*, Tenet®\*, Revystar®, Verdict®\* 520, and Versys® without loss of efficacy or adverse crop effects.

Seek advice from BASF Australia Ltd before using products other than Liberty® Ultra Herbicide or those mentioned above on LibertyLink® Canola varieties.

#### Fallow

Liberty® Ultra Herbicide is compatible with most residual herbicides e.g. simazine, diuron, oxyfluorfen (Goal®), norfluzuron (Solicam®) and oryzalin (Surflan®), and with glyphosate, metsulfuron-methyl, paraquat, Sharpen® and Voraxor®.

### RECROPPING INTERVALS

**Minimum recropping intervals apply for crops following Liberty® Ultra Herbicide application.**

For advice on crops not listed below, contact the manufacturer, BASF Australia Ltd.

Crop	Minimum recropping interval
Cereals, pulses, oilseeds	1 day

### RESISTANT WEEDS WARNING

GROUP 10 HERBICIDE

Liberty® Ultra Herbicide is a member of the phosphinic acid group of herbicides. Liberty® Ultra Herbicide is an inhibitor of glutamine synthetase. For weed resistance management Liberty® Ultra Herbicide is a Group 10 herbicide. Some naturally-occurring weed biotypes resistant to Liberty® Ultra Herbicide and other Group 10 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Liberty or other Group 10 herbicides. Since occurrence of resistant weeds is difficult to detect prior to use, BASF Australia Ltd accepts no liability for any losses that may result from the failure of Liberty® Ultra Herbicide to control resistant weeds.

Where resistance to this product is suspected this should be reported either directly to the Australian Pesticides and Veterinary Medicines Authority (APVMA) or through BASF Australia Ltd.

## **LibertyLink® Canola varieties (with tolerance to Liberty® Ultra Herbicide) Canola Resistance and Crop Management Plans**

Detailed management plans have been developed that outline sound agronomic practices and integrated weed management principles, designed to optimise the performance of LibertyLink® Canola varieties (with tolerance to Liberty® Ultra Herbicide), minimise the potential for the development of herbicide resistance in weed populations and ensure compliance with the Conditions of the LibertyLink® Canola OGTR approval.

The LibertyLink® Canola varieties (with tolerance to Liberty® Ultra Herbicide) Resistance Management and Crop Management Plans should be consulted prior to the use of this product on LibertyLink® Canola varieties. Implementation of the management plans are an essential part of herbicide resistance management. This product is supplied by BASF Australia Ltd on condition that it is used in conjunction with the LibertyLink® Canola varieties (with tolerance to Liberty® Ultra Herbicide) Resistance Management and Crop Management Plans (available on [www.crop-solutions.basf.com.au](http://www.crop-solutions.basf.com.au)).

Pre-plant paddock preparation: Control all existing weeds by cultivation or by using a sufficient rate of knockdown herbicide. This may be a glyphosate or paraquat/diquat-based product.

### **PRECAUTION**

#### **Re-entry period**

DO NOT enter treated areas until spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Very toxic to aquatic life. DO NOT contaminate streams, rivers or waterways with this product or the used container.

### **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

Do not apply on desirable foliage (other than LibertyLink® Canola varieties with tolerance to Liberty® Ultra Herbicide) or allow drift onto the foliage or green bark of trees and vines, as damage will occur.

### **INTEGRATED PEST MANAGEMENT**

Toxic to beneficial arthropods. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas.

### **STORAGE**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

### **DISPOSAL**

Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging or unused product to an approved waste management facility. If an approved waste management facility is not available, dispose of empty container or unused product in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

#### **110 L container**

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the container to the point of purchase.

#### **1000 L container**

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty product as required into application equipment. Do not attempt to breach the valve system or filling point, or contaminate the container with water or other products. Ensure that equipment used in transfer of the product is disconnected, triple rinsed

and drained after each use. When the container is empty, close all caps and valves and return the container to the point of purchase.

### **SAFETY DIRECTIONS**

Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. When using together with other products, consult their label safety directions. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow length chemical resistant gloves and goggles. When using the prepared spray wear cotton overalls, over normal clothing, buttoned to the neck and wrist, elbow length chemical resistant gloves and a half facepiece respirator. Wash hands after use. After each day's use, wash gloves, goggles, respirator and contaminated clothing.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet available from your supplier.

### **CONDITIONS OF SALE**

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Competition and Consumer Act 2010 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 94738/143503

Batch No:

Date of Manufacture:

LibertyLink® and Liberty® are registered trademarks of BASF.

© BASF 2026

\* Other registered trademarks

### **BASF Australia Ltd**

ABN 62 008 437 867

Level 23, 40 City Road

Southbank VICTORIA 3006

[crop-solutions.basf.com.au](http://crop-solutions.basf.com.au)

**FOR SPECIALIST ADVICE IN AN  
EMERGENCY ONLY  
PHONE 1800 803 440  
TOLL FREE - ALL HOURS - AUSTRALIA WIDE**

**GHS STATEMENTS**

Causes serious eye damage. Causes skin irritation. Harmful if inhaled. May be harmful if swallowed. May cause damage to organs through prolonged or repeated exposure. May damage fertility. Suspected of damaging the unborn child. Very toxic to aquatic life. Very toxic to aquatic life with long lasting effects. If medical advice is needed, have product container or label at hand. Keep out of reach of children. Read carefully and follow all instructions. Use only outdoors or in a well-ventilated area. Wear protective gloves, protective clothing and eye protection or face protection. Do not breathe dust/gas/mist/vapours. Do not handle until all safety precautions have been read and understood. Wash contaminated body parts thoroughly after handling. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or physician. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF exposed or concerned: Get medical attention. IF ON SKIN: Wash with plenty of soap and water. Collect spillage. Take off contaminated clothing and wash it before reuse. Store locked up.